


# Ecosystem Restoration on the South Farallon Islands



# Importance of islands



Farallon arboreal salamander



Harbor seals


- Endemic species
- 3% of earth's surface, but **15-20%** of birds, reptiles and plants
- Critical habitat

# Most extinctions are on islands



## Extinctions since 1600

- **94%** of extinct birds
- **90%** of extinct reptiles
- **65%** of extinct mammals
- **68%** of extinct plants




# Cause of island animal extinctions:



**Invasive species (55-67%)**

# House mice on islands





**Farallon Islands**

# Ashy storm-petrels:

*Limited and rare*

- Limited breeding grounds
  1. Farallon Is.: **50% of population**
  2. Channel Is. (Southern CA)
- Rare (5,000-10,000 birds)



# Ashy storm-petrels:

## *Characteristics of a vulnerable species*

- Long-lived
- Slow reproduction
- Months of care



# Ashy storm-petrels in decline

- **42% Farallons population decline** (breeding birds)
- **Identified as a species at risk**
  - IUCN
  - Audubon



# Ashy storm-petrels: *Predators*

- Gulls (natural predator)
- Burrowing owls  
(artificially supported by  
mice)
- House mice (rarely  
documented)






## **Mice & ashy storm-petrels:**


*The hyperpredation link*



# Hyperpredation



No mice



Fall

Winter

Spring

# Why remove mice?

- Ashy storm-petrels (burrowing owls leave)
- Other seabirds (disturbance)
- Native insects
- Native plants

# Mouse eradication:

*100% removal*

- *Introduced predators*
- *Trapping*
- *Virus*
- **Rodenticide bait**



# Bait delivery options

## Bait stations




- Get every mouse
- Minimize disturbance
- Ecosystem benefits must outweigh risks

## Aerial broadcast



# Timing

Farallon mouse population cycle (mouse trap success by month)



 **Best time for eradication**

 **Not ideal for eradication**

# Issues to consider

- Ecosystem disturbance?

- Effects of rodenticide?



# Rodenticide use

HAZARD =

1) Toxicity + 2) Exposure

# Rodenticides for conservation:

## *Toxicity risk*

- Not affected:
  - Invertebrates
  - Plants
- Affected:
  - Vertebrates (e.g. mice)



# Rodenticides for conservation:

## *Minimizing exposure risk*

- Bait presentation
  - *Bait stations* inaccessible by most birds
  - *Grain-based pellet* unattractive to pinnipeds, nearly all seabirds, insect-eating songbirds, most fish
  - *Green coloration* unattractive to many grain-eating songbirds
- Dosage (body size)
  - No risk to pinnipeds (or humans)



# **Rodenticides:**

*Minimizing exposure risk*

- Seasonal timing
- Non-target capture/hold

# Rodenticides:

## *Aerial broadcast considerations*

- Precision application
- Bait pellets disintegrate quickly
- Rodenticide doesn't dissolve into water
- Low rodenticide concentration



# Compliance

- National Environmental Policy Act (NEPA)
- Clean Air Act (CAA)
- Clean Water Act (CWA)
- Endangered Species Act (ESA)
- National Historic Preservation Act (NHPA)
- Wilderness Act (WA)
- Marine Mammal Protection Act (MMPA)
- Migratory Bird Treaty Act (MBTA)
- National Marine Sanctuaries Act (NMSA)
- Coastal Zone Management Act (CZMA)
- Federal Insecticide Fungicide and Rodenticide Act (FIFRA)

# Partners



Farallon National Wildlife Refuge (USFWS)



Island Conservation



PRBO Conservation Science



National Fish & Wildlife Foundation



*S.S. Luckenbach* Trustee Council

**We want your input!**



**Jesse Irwin / Joelle Buffa**  
**SF Bay National Wildlife Refuge Complex**  
**PO Box 524, Newark CA 94560**  
**(510) 792-0222**  
[sfbaynwrc@fws.gov](mailto:sfbaynwrc@fws.gov)  
**(510) 792-5828 fax**